

Hypersurface of a Finsler space subjected to a Kropina change with an h -vector

M. K. GUPTA*

Department of Pure & Applied Mathematics
Guru Ghasidas Vishwavidyalaya
Bilaspur (C.G.), India

P. N. PANDEY

Department of Mathematics
University of Allahabad
Allahabad, India

Abstract

The concept of h -vector was introduced by H. Izumi in 1980. Recently we have obtained the Cartan connection for the Finsler space whose metric is given by Kropina change with an h -vector. In 1985, M. Matsumoto studied the theory of Finsler hypersurface. In this paper, we derive certain geometrical properties of a Finslerian hypersurface subjected to a Kropina change with an h -vector.

Keywords: Finsler space, hypersurface, Kropina change, h -vector.

2000 Mathematics Subject Classification: **53B40**.

1 Introduction

In 1984, C. Shibata [7]¹ dealt with a change of Finsler metric which is called a β -change of metric. A remarkable class of β -change is Kropina change ${}^*L(x, y) = \frac{L^2(x, y)}{b_i(x)y^i}$. If $L(x, y)$ is a metric function of a Riemannian space then ${}^*L(x, y)$ reduces to the metric function of a Kropina space. Such a Finsler metric was introduced by V.K. Kropina [4].

*Corresponding author. E-mail: mkgiaps@gmail.com

¹Numbers in square brackets refer to the references at the end of the paper.

H. Izumi [3] while studying the conformal transformation of Finsler spaces, introduced the h -vector b_i , which is v -covariant constant with respect to the Cartan connection and satisfies $L C_{ij}^h b_h = \rho h_{ij}$, $\rho \neq 0$. Thus if b_i is an h -vector then

$$(1.1) \quad (i) \ b_i|_k = 0, \quad (ii) \ L C_{ij}^h b_h = \rho h_{ij}.$$

This gives

$$(1.2) \quad L \dot{\partial}_j b_i = \rho h_{ij}.$$

Since $\rho \neq 0$ and $h_{ij} \neq 0$ for $h_{ij} = 0$ implies $n = 1$ which is not true, the h -vector b_i depends not only on positional coordinates but also on directional arguments. Izumi [3] showed that ρ is independent of directional arguments. The present authors [2] obtain the relation between the Cartan connections of $F^n = (M^n, L)$ and ${}^*F^n = (M^n, {}^*L)$ where ${}^*L(x, y)$ is obtained by the transformation

$$(1.3) \quad {}^*L(x, y) = \frac{L^2(x, y)}{b_i(x, y) y^i}$$

and $b_i(x, y)$ is an h -vector in (M^n, L) .

M. Matsumoto [5] presented a systematic theory of Finslerian hypersurface. The present authors [1] obtained certain results for the Finslerian hypersurface.

In this paper, certain geometrical properties of a Finslerian hypersurface subjected to a Kropina change with an h -vector, were discussed.

The terminologies and notations are referred to Matsumoto [6].

2 Preliminaries

Let M^n be an n -dimensional smooth manifold and $F^n = (M^n, L)$ be an n -dimensional Finsler space equipped with a metric function $L(x, y)$ on M^n . The normalized supporting element, the metric tensor, the angular metric tensor and Cartan tensor are defined by $l_i = \dot{\partial}_i L$, $g_{ij} = \frac{1}{2} \dot{\partial}_i \dot{\partial}_j L^2$, $h_{ij} = L \dot{\partial}_i \dot{\partial}_j L$ and $C_{ijk} = \frac{1}{2} \dot{\partial}_k g_{ij}$ respectively. Throughout this paper, we use the symbols $\dot{\partial}_i$ and ∂_i for $\partial/\partial y^i$ and $\partial/\partial x^i$ respectively. The Cartan

connection in F^n is given as $CT = (F_{jk}^i, G_j^i, C_{jk}^i)$. The h - and v -covariant derivatives of a covariant vector $X_i(x, y)$ with respect to the Cartan connection are given by

$$(2.1) \quad X_{i|j} = \partial_j X_i - (\dot{\partial}_h X_i) G_j^h - F_{ij}^r X_r,$$

and

$$(2.2) \quad X_i|_j = \dot{\partial}_j X_i - C_{ij}^r X_r.$$

A hypersurface M^{n-1} of the underlying manifold M^n may be represented parametrically by the equations $x^i = x^i(u^\alpha)$, where u^α are the Gaussian coordinates on M^{n-1} (Latin indices run from 1 to n , while Greek indices take values from 1 to $n-1$). We assume that the matrix of projection factors $B_\alpha^i = \partial x^i / \partial u^\alpha$ is of rank $n-1$. If the supporting element y^i at a point $u = (u^\alpha)$ of M^{n-1} is assumed to be tangent to M^{n-1} , we may then write $y^i = B_\alpha^i(u) v^\alpha$ so that $v = (v^\alpha)$ is thought of as the supporting element of M^{n-1} at the point u^α . Since the function $\underline{L}(u, v) = L(x(u), y(u, v))$ gives rise to a Finsler metric on M^{n-1} , we get an $(n-1)$ -dimensional Finsler space $F^{n-1} = (M^{n-1}, \underline{L}(u, v))$.

At each point u^α of F^{n-1} , a unit normal vector $N^i(u, v)$ is defined by

$$(2.3) \quad g_{ij} B_\alpha^i N^j = 0, \quad g_{ij} N^i N^j = 1.$$

The inverse projection factors $B_i^\alpha(u, v)$ of B_α^i are defined as

$$(2.4) \quad B_i^\alpha = g^{\alpha\beta} g_{ij} B_\beta^j,$$

where $g^{\alpha\beta}$ is the inverse of the metric tensor $g_{\alpha\beta}$ of F^{n-1} .

From (2.3) and (2.4), it follows that

$$(2.5) \quad B_\alpha^i B_i^\beta = \delta_\alpha^\beta, \quad B_\alpha^i N_i = 0, \quad N^i B_i^\alpha = 0, \quad N^i N_i = 1,$$

and further

$$(2.6) \quad B_\alpha^i B_j^\alpha + N^i N_j = \delta_j^i.$$

For the induced Cartan connection $ICT = (F_{\beta\gamma}^\alpha, G_\beta^\alpha, C_{\beta\gamma}^\alpha)$ on F^{n-1} , the second fundamental h -tensor $H_{\alpha\beta}$ and the normal curvature vector H_α are given by

$$(2.7) \quad H_{\alpha\beta} = N_i (B_{\alpha\beta}^i + F_{jk}^i B_\alpha^j B_\beta^k) + M_\alpha H_\beta,$$

and

$$(2.8) \quad H_\alpha = N_i (B_{0\alpha}^i + G_j^i B_\alpha^j),$$

where $M_\alpha = C_{ijk} B_\alpha^i N^j N^k$, $B_{\alpha\beta}^i = \partial^2 x^i / \partial u^\alpha \partial u^\beta$ and $B_{0\alpha}^i = B_{\beta\alpha}^i v^\beta$.

The equations (2.7) and (2.8) yield

$$(2.9) \quad H_{0\alpha} = H_{\beta\alpha} v^\beta = H_\alpha, \quad H_{\alpha 0} = H_{\alpha\beta} v^\beta = H_\alpha + M_\alpha H_0.$$

The second fundamental v-tensor $M_{\alpha\beta}$ is defined as:

$$(2.10) \quad M_{\alpha\beta} = C_{ijk} B_\alpha^i B_\beta^j N^k.$$

The relative h - and v -covariant derivatives of B_α^i and N^i are given by

$$(2.11) \quad \begin{aligned} B_{\alpha|\beta}^i &= H_{\alpha\beta} N^i, & B_\alpha^i|_\beta &= M_{\alpha\beta} N^i, \\ N_{|\beta}^i &= -H_{\alpha\beta} B_j^\alpha g^{ij}, & N^i|_\beta &= -M_{\alpha\beta} B_j^\alpha g^{ij}. \end{aligned}$$

Let $X_i(x, y)$ be a vector field of F^n . The relative h - and v -covariant derivatives of X_i are given by

$$(2.12) \quad X_{i|\beta} = X_{i|j} B_\beta^j + X_i|_j N^j H_\beta, \quad X_i|_\beta = X_i|_j B_\beta^j.$$

Matsumoto [5] defined different kinds of hyperplanes and obtained their characteristic conditions, which are given in the following lemmas:

Lemma 2.1. *A hypersurface F^{n-1} is a hyperplane of the first kind if and only if $H_\alpha = 0$ or equivalently $H_0 = 0$.*

Lemma 2.2. *A hypersurface F^{n-1} is a hyperplane of the second kind if and only if $H_{\alpha\beta} = 0$.*

Lemma 2.3. *A hypersurface F^{n-1} is a hyperplane of the third kind if and only if $H_{\alpha\beta} = 0 = M_{\alpha\beta}$.*

3 The Finsler space ${}^*F^n = (M^n, {}^*L)$

If we denote $b_i y^i$ by β then indicatory property of h_{ij} yield $\dot{\partial}_i \beta = b_i$. Throughout this paper, the geometric objects associated with ${}^*F^n$ will be asterisked. We shall use

the notation $L_{ij} = \dot{\partial}_i \dot{\partial}_j L$, $L_{ijk} = \dot{\partial}_k L_{ij}$, ... etc. From (1.3), we get

$$(3.1) \quad {}^*L_{ij} = (2\tau - \rho\tau^2) L_{ij} + \frac{2\tau^2}{\beta} m_i m_j,$$

$$(3.2) \quad \begin{aligned} {}^*L_{ijk} = & (2\tau - \rho\tau^2) L_{ijk} + \frac{2\tau}{\beta} (\rho\tau - 1) (m_i L_{jk} + m_j L_{ik} + m_k L_{ij}) \\ & - \frac{2\tau^2}{L\beta} (m_i m_j l_k + m_j m_k l_i + m_k m_i l_j) - \frac{6\tau^2}{\beta^2} m_i m_j m_k, \end{aligned}$$

where $\tau = L/\beta$, $m_i = b_i - \frac{1}{\tau} l_i$. The normalized supporting element, the metric tensor of ${}^*F^n$ are obtained as [2]

$$(3.3) \quad {}^*l_i = 2\tau l_i - \tau^2 b_i,$$

$$(3.4) \quad {}^*g_{ij} = (2\tau^2 - \rho\tau^3) g_{ij} + 3\tau^4 b_i b_j - 4\tau^3 (l_i b_j + b_i l_j) + (4\tau^2 + \rho\tau^3) l_i l_j,$$

Differentiating the angular metric tensor h_{ij} with respect to y^k , we get

$$\dot{\partial}_h h_{ij} = 2C_{ijk} - \frac{1}{L} (l_i h_{jk} + l_j h_{ik}),$$

which gives

$$(3.5) \quad L_{ijk} = \frac{2}{L} C_{ijk} - \frac{1}{L^2} (h_{ij} l_k + h_{jk} l_i + h_{ki} l_j).$$

Taking (3.5) into account, (3.2) can be rewritten as

$$(3.6) \quad {}^*C_{ijk} = (2\tau^2 - \rho\tau^3) C_{ijk} - \frac{\tau^2}{2\beta} (4 - 3\rho\tau) (h_{ij} m_k + h_{jk} m_i + h_{ki} m_j) - \frac{6\tau^2}{\beta} m_i m_j m_k.$$

The inverse metric tensor of ${}^*F^n$ is derived as follows [2]:

$$(3.7) \quad \begin{aligned} {}^*g^{ij} = & (2\tau^2 - \rho\tau^3)^{-1} \left[g^{ij} - \frac{2\tau}{2b^2\tau - \rho} b^i b^j + \frac{4 - \rho\tau}{2b^2\tau - \rho} (l^i b^j + b^i l^j) \right. \\ & \left. - \frac{3\rho b^2\tau^3 - \rho^2\tau^2 - 4b^2\tau^2 - 2\rho\tau + 8}{\tau(2b^2\tau - \rho)} l^i l^j \right] \end{aligned}$$

where b is the magnitude of the vector $b^i = g^{ij} b_j$.

We obtained the relation between the Cartan connection coefficients F_{jk}^i and ${}^*F_{jk}^i$ as [2]

$$(3.8) \quad {}^*F_{jk}^i = F_{jk}^i + D_{jk}^i.$$

Transvecting by y^j and using $F_{jk}^i y^j = G_k^i$, we get

$$(3.9) \quad {}^*G_k^i = G_k^i + D_{0k}^i,$$

where the subscript '0' denotes the contraction by the supporting element y^j . Further transvecting (3.9) by y^k and using $G_k^i y^k = 2G^i$, we get

$$(3.10) \quad 2{}^*G^i = 2G^i + D_{00}^i.$$

Differentiating (3.9) partially with respect to y^h and using $\dot{\partial}_h G_k^i = G_{kh}^i$, we have

$$(3.11) \quad {}^*G_{kh}^i = G_{kh}^i + \dot{\partial}_h D_{0k}^i,$$

where G_{kh}^i are the Berwald connection coefficients. The expressions of D_{00}^i , D_{0j}^i and D_{jk}^i are given by

$$(3.12) \quad D_{00}^i = l^i \tau (A - E_{00}) - \frac{2\tau^2}{2 - \rho\tau} A m^i + \frac{2L\tau}{2 - \rho\tau} \left(\frac{1}{\beta} \beta_0 m^i - F_0^i \right),$$

$$(3.13) \quad D_{0j}^i = l^i \left\{ \frac{1}{\tau} \left(\frac{2}{\beta} b^2 - \frac{\rho}{L} \right)^{-1} G_{\beta j} + \frac{1}{\tau} G_j \right\} - \frac{2m^i}{2 - \rho\tau} \left(\frac{2}{\beta} b^2 - \frac{\rho}{L} \right)^{-1} G_{\beta j} + \frac{L}{2\tau - \rho\tau^2} G_j^i,$$

$$(3.14) \quad D_{ik}^j = l^j \left\{ \frac{1}{\tau} \left(\frac{2}{\beta} b^2 - \frac{\rho}{L} \right)^{-1} H_{\beta ik} + \frac{1}{\tau} H_{ik} \right\} - \frac{2m^j}{2 - \rho\tau} \left(\frac{2}{\beta} b^2 - \frac{\rho}{L} \right)^{-1} H_{\beta ik} + \frac{L}{2\tau - \rho\tau^2} H_{ik}^j,$$

where

$$(3.15) \quad \begin{aligned} G_{ij} = & \frac{\tau^2}{\beta} (m_i \beta_j - m_j \beta_i) - \tau^2 F_{ij} - \frac{1}{2} (2\tau - \rho\tau^2) L_{ijr} D_{00}^r - \frac{\tau^2}{2} \rho_0 L_{ij} \\ & - \frac{\tau}{\beta} D_{00}^r \sum_{ijr} m_i \left\{ (\rho\tau - 1) L_{jr} - \frac{\tau}{\beta} m_j b_r \right\} + \frac{\tau}{\beta} \beta_0 \left((\rho\tau - 1) L_{ij} - \frac{3\tau}{\beta} m_i m_j \right), \end{aligned}$$

$$(3.16) \quad G_j = \tau^2 (F_{j0} - E_{j0}),$$

$$(3.17) \quad \begin{aligned} H_{jik} = & \frac{(\rho\tau^2 - 2\tau)}{2} \{ L_{ijr} D_{0k}^r + L_{jkr} D_{0i}^r - L_{kir} D_{0j}^r \} \\ & - \frac{\tau}{\beta} D_{0k}^r \sum_{ijr} m_i \left\{ (\rho\tau - 1) L_{jr} - \frac{\tau}{\beta} m_j b_r \right\} - \frac{\tau}{\beta} D_{0i}^r \sum_{jkr} m_j \left\{ (\rho\tau - 1) L_{kr} - \frac{\tau}{\beta} m_k b_r \right\} \\ & + \frac{\tau}{\beta} D_{0j}^r \sum_{kir} m_k \left\{ (\rho\tau - 1) L_{ir} - \frac{\tau}{\beta} m_i b_r \right\} - \frac{\tau^2}{2} (\rho_k L_{ij} + \rho_i L_{jk} - \rho_j L_{ki}) \\ & + \frac{\tau}{\beta} \left\{ \beta_k ((\rho\tau - 1) L_{ij} - \frac{3\tau}{\beta} m_i m_j) + \beta_i ((\rho\tau - 1) L_{jk} - \frac{3\tau}{\beta} m_j m_k) \right. \\ & \left. - \beta_j ((\rho\tau - 1) L_{ki} - \frac{3\tau}{\beta} m_k m_i) \right\}, \end{aligned}$$

$$(3.18) \quad H_{ik} = \frac{\tau^2}{\beta}(m_i\beta_k + m_k\beta_i) - \tau^2 E_{ik} - \frac{1}{2}(G_{ik} + G_{ki}),$$

and

$$(3.19) \quad A = \left(\frac{2}{\beta} \beta_0 m^2 - 2F_{\beta 0} \right) \left(\frac{2}{\beta} b^2 - \frac{\rho}{L} \right)^{-1},$$

$$F_j^i = g^{ir} F_{rj}, \quad G_j^i = g^{ik} G_{kj}, \quad H_{ik}^j = g^{jm} H_{mik},$$

$$2E_{ij} = b_{i|j} + b_{j|i}, \quad 2F_{ij} = b_{i|j} - b_{j|i}, \quad \beta_j = \beta_{|j}, \quad \rho_k = \rho_{|k} = \partial_k \rho.$$

The symbol $\sum_{ijk}$ denote cyclic interchange of indices i, j, k and summation.

Lemma 3.1. [2] *If the h-vector b_i is gradient then the scalar ρ is constant.*

Lemma 3.2. [2] *For the Kropina change with h-vector, the difference tensor D_{jk}^i vanishes if and only if the vector b_i is parallel with respect to the Cartan connection of F^n , and then the Berwald connection coefficients for both the spaces F^n and ${}^*F^n$ are the same.*

4 Hypersurface ${}^*F^{n-1}$ of the space ${}^*F^n$

Let us consider a Finslerian hypersurface $F^{n-1} = (M^{n-1}, \underline{L}(u, v))$ of F^n and a Finslerian hypersurface ${}^*F^{n-1} = (M^{n-1}, {}^*\underline{L}(u, v))$ of ${}^*F^n$. Let N^i be the unit normal vector at a point of F^{n-1} . The functions $B_\alpha^i(u)$ may be considered as the components of $n-1$ linearly independent vectors tangent to F^{n-1} and they are invariant under the Randers conformal change. Then the unit normal vector ${}^*N^i(u, v)$ of ${}^*F^{n-1}$ is uniquely determined by

$$(4.1) \quad {}^*g_{ij} B_\alpha^i {}^*N^j = 0, \quad {}^*g_{ij} {}^*N^i {}^*N^j = 1.$$

The inverse projection factors ${}^*B_i^\alpha$ are uniquely defined along ${}^*F^{n-1}$ by

$$(4.2) \quad {}^*B_i^\alpha = {}^*g^{\alpha\beta} {}^*g_{ij} B_\beta^j,$$

where ${}^*g^{\alpha\beta}$ is the inverse of the metric tensor ${}^*g_{\alpha\beta}$ of ${}^*F^{n-1}$.

From (4.2), it follows that

$$(4.3) \quad B_\alpha^i {}^*B_i^\beta = \delta_\alpha^\beta, \quad B_\alpha^i {}^*N_i = 0, \quad {}^*N^i {}^*B_i^\alpha = 0, \quad {}^*N^i {}^*N_i = 1.$$

Transvecting (2.3) by v^α , we get

$$(4.4) \quad y_j N^j = 0.$$

Transvecting (3.4) by $N^i N^j$ and paying attention to (2.3) and (4.4), we have

$$(4.5) \quad {}^*g_{ij} N^i N^j = (2\tau^2 - \rho\tau^3) + 3\tau^4 (b_i N^i)^2,$$

which shows that $N^j / \sqrt{(2\tau^2 - \rho\tau^3) + 3\tau^4 (b_i N^i)^2}$ is a unit vector. Again transvecting (3.4) by $B_\alpha^i N^j$ and using (2.3) and (4.4), we get

$$(4.6) \quad {}^*g_{ij} B_\alpha^i N^j = (b_j N^j)(3\tau^4 b_i B_\alpha^i - 4\tau^3 l_i B_\alpha^i).$$

This shows that the vector N^j is normal to ${}^*F^{n-1}$ if and only if

$$(b_j N^j)(3\tau^4 b_i B_\alpha^i - 4\tau^3 l_i B_\alpha^i) = 0.$$

This implies at least one of the following :

$$(a) \quad 3\tau^4 b_i B_\alpha^i - 4\tau^3 l_i B_\alpha^i = 0, \quad (b) \quad b_j N^j = 0.$$

The transvection of $3\tau^4 b_i B_\alpha^i - 4\tau^3 l_i B_\alpha^i = 0$ by v^α gives $3\tau^4 b_i y^i - 4\tau^3 l_i y^i = 0$, i.e. $\tau = L/\beta = 0$, which is not possible. Therefore $3\tau^4 b_i B_\alpha^i - 4\tau^3 l_i B_\alpha^i \neq 0$. Hence

$$(4.7) \quad b_j N^j = 0.$$

This shows that the vector N^j is normal to ${}^*F^{n-1}$ if and only if b_j is tangent to F^{n-1} .

From (4.5), (4.6) and (4.7), we can say that $N^j / \sqrt{(2\tau^2 - \rho\tau^3)}$ is a unit normal vector of ${}^*F^{n-1}$. Therefore in view of (4.1), we get

$$(4.8) \quad {}^*N^i = N^i / \sqrt{(2\tau^2 - \rho\tau^3)},$$

which, in view of (3.4), (4.4) and (4.7), gives

$$(4.9) \quad {}^*N_i = {}^*g_{ij} {}^*N^j = \sqrt{(2\tau^2 - \rho\tau^3)} N_i.$$

Thus, we have:

Theorem 4.1. *Let ${}^*F^n$ be the Finsler space obtained from F^n by a Kropina change (1.3) with h -vector. If ${}^*F^{n-1}$ and F^{n-1} are the hypersurfaces of these spaces then the vector b_i is tangential to the hypersurface F^{n-1} if and only if every vector normal to F^{n-1} is also normal to ${}^*F^{n-1}$.*

In view of (4.4) and (4.7), the vector m_i appearing in (3.2) satisfies

$$(4.10) \quad m_i N^i = 0.$$

As $h_{ij} = g_{ij} - l_i l_j$, equations (2.3) and (4.4) yield

$$(4.11) \quad h_{ij} B_\alpha^i N^i = 0.$$

Transvecting (3.6) by $B_\alpha^i B_\beta^j N^k$ and using (4.10) and (4.11), we get

$$(4.12) \quad {}^*C_{ijk} B_\alpha^i B_\beta^j N^k = (2\tau^2 - \rho\tau^3) C_{ijk} B_\alpha^i B_\beta^j N^k.$$

Using (2.10) and (4.8), equation (4.12) may be written as

$$(4.13) \quad {}^*M_{\alpha\beta} = \sqrt{(2\tau^2 - \rho\tau^3)} M_{\alpha\beta}.$$

From (2.8), (3.13) and (4.9), we get

$${}^*H_\alpha = \sqrt{(2\tau^2 - \rho\tau^3)} (H_\alpha + N_i D_{0k}^i B_\alpha^k).$$

Transvecting by v^α and using $v^\alpha B_\alpha^k = y^k$, we get

$$(4.14) \quad {}^*H_0 = \sqrt{(2\tau^2 - \rho\tau^3)} (H_0 + N_i D_{00}^i).$$

Transvecting (3.12) by N_i and using (4.4) and (4.10), we get

$$(4.15) \quad D_{00}^i N_i = -\frac{2L\tau}{2 - \rho\tau} F_0^i N_i.$$

If the vector b_i is gradient, i.e. $b_{i|j} = b_{j|i}$, then

$$(4.16) \quad F_{ij} = 0,$$

and then by Lemma 3.1, we have

$$(4.17) \quad \rho_i = 0.$$

Therefore (4.15) becomes $D_{00}^i N_i = 0$, and then equation (4.14) reduces to

$${}^*H_0 = \sqrt{(2\tau^2 - \rho\tau^3)}H_0.$$

Thus, in view of Lemma 2.1, we have:

Theorem 4.2. *Let the h -vector $b_i(x, y)$ be a gradient and tangent to the hypersurface F^{n-1} . Then the hypersurface F^{n-1} is a hyperplane of the first kind if and only if the hypersurface ${}^*F^{n-1}$ is a hyperplane of the first kind.*

Taking the relative h -covariant differentiation of (4.7) with respect to the Cartan connection of F^{n-1} , we get

$$b_{i|\beta} N^i + b_i N^i_{|\beta} = 0.$$

Using (2.11) and (2.12), the above equation gives

$$(b_{i|j} B_\beta^j + b_i|_j N^j H_\beta) N^i - b_i H_{\alpha\beta} B_j^\alpha g^{ij} = 0.$$

Transvecting by v^β and using (2.9), we get

$$(4.18) \quad b_{i|0} N^i = (H_\alpha + M_\alpha H_0) B_j^\alpha b^j - b_i|_j H_0 N^i N^j.$$

For a hypersurface of the first kind, $H_0 = 0 = H_\alpha$. Then (4.18) reduces to $b_{i|0} N^i = 0$.

If the vector b_i is gradient, i.e. $b_{i|j} = b_{j|i}$, then we get

$$(4.19) \quad E_{i0} N^i = b_{i|0} N^i = \beta_i N^i = 0.$$

Therefore (3.16) gives

$$(4.20) \quad G_j N^j = 0.$$

Transvecting (3.15) by $b^i N^j$ and using (4.10), (4.16), (4.17) and (4.19), we get

$$(4.21) \quad G_{ij} b^i N^j = G_{\beta j} N^j = 0.$$

Again transvecting (3.15) by $N^i B_\alpha^j$ and using $D_{00}^i N_i = 0$ and $b_i N^i = 0$, we obtain

$$(4.22) \quad G_{ij} N^i B_\alpha^j = 0.$$

Transvecting (3.13) by $N_i B_\alpha^j$ and using (4.4), (4.10) and (4.22), we get

$$(4.23) \quad D_{0j}^i N_i B_\alpha^j = 0.$$

Transvecting (3.12) by L_{ijk} and using (3.5) and (4.16), we have

$$(4.24) \quad L_{ijk} D_{00}^i = (A - \beta_0) \left[- \left(\frac{\tau}{L^2} + \frac{4\rho\tau^2}{L^2(2 - \rho\tau)} \right) h_{jk} + \frac{2\tau^2}{L^2(2 - \rho\tau)} (m_i l_k + m_k l_i) \right].$$

Transvecting (3.13) by $N^j B_\alpha^k h_{ki}$ and using (4.10) and (4.24), we get

$$(4.25) \quad D_{0j}^i N^j B_\alpha^k h_{ik} = 0.$$

Again transvecting (3.13) by $b_i N^j$ and using (4.20) and (4.21), we get

$$(4.26) \quad D_{0j}^i b_i N^j = 0.$$

Transvecting (3.17) by $N^j B_\alpha^i B_\beta^k$ and using (4.10), (4.23), (4.25) and (4.26), we obtain

$$(4.27) \quad H_{jik} N^j B_\alpha^i B_\beta^k = \frac{(\rho\tau^2 - 2\tau)}{2} N^j B_\alpha^i B_\beta^k (L_{ijr} D_{0k}^r + L_{jkr} D_{0i}^r - L_{kir} D_{0j}^r).$$

Let the h -vector b_i satisfies the condition

$$(4.28) \quad b_{r|0} C_{ij}^r = 0,$$

then

$$(4.29) \quad \beta_r C_{ij}^r = 0.$$

Transvecting G_{rk} by $C_{ij}^r N^j B_\alpha^i B_\beta^k$ and using (4.16), (4.17), (4.24) and (4.29), we get

$$\begin{aligned} C_{ij}^r N^j B_\alpha^i B_\beta^k G_{rk} &= -C_{ij}^r N^j B_\alpha^i B_\beta^k \left[(A - \beta_0) \left(\frac{\tau}{L^2} + \frac{4\rho\tau^2}{L^2(2 - \rho\tau)} \right) h_{rk} \right. \\ &\quad \left. + \frac{\tau}{\beta} (\rho\tau - 1) D_{00}^s m_s L_{rk} - \frac{\tau}{\beta} \beta_0 (\rho\tau - 1) L_{rk} \right]. \end{aligned}$$

Using (2.10), above equation is simplified as

$$C_{ij}^r N^j B_\alpha^i B_\beta^k G_{rk} = \lambda M_{\alpha\beta},$$

where we put

$$\lambda = -(A - \beta_0) \left(\frac{\tau}{L^2} + \frac{4\rho\tau^2}{L^2(2 - \rho\tau)} \right) + (A - \beta_0)(\rho\tau - 1) \frac{2\tau^3 m^2}{L\beta(2 - \rho\tau)} + \frac{\tau}{L\beta} \beta_0(\rho\tau - 1).$$

Hence we get

$$(4.30) \quad L_{ijr} N^j B_\alpha^i B_\beta^k D_{0k}^r = \frac{2\lambda}{2\tau - \rho\tau^2} M_{\alpha\beta}.$$

By interchanging the indices i and k , we have

$$(4.31) \quad L_{jkr} N^j B_\alpha^i B_\beta^k D_{0i}^r = \frac{2\lambda}{2\tau - \rho\tau^2} M_{\alpha\beta}.$$

Transvecting (3.15) by N^j and using (4.10), (4.16), (4.19) and (4.24), we get

$$G_{ij} N^j = \mu N_i,$$

where we put

$$\mu = \frac{1}{2}(2\tau - \rho\tau^2)(A - \beta_0) \left(\frac{\tau}{L^2} + \frac{4\rho\tau^2}{L^2(2 - \rho\tau)} \right) - \frac{\tau}{L\beta} m_s D_{00}^s + \frac{\tau}{L\beta} \beta_0(\rho\tau - 1).$$

Using this, we obtain

$$(4.32) \quad L_{kir} N^j B_\alpha^i B_\beta^k D_{0j}^r = \frac{2\mu}{2\tau - \rho\tau^2} M_{\alpha\beta}.$$

Using (4.30), (4.31), (4.32) in (4.27), we get

$$(4.33) \quad H_{jik} N^j B_\alpha^i B_\beta^k = (\mu - 2\lambda) M_{\alpha\beta}.$$

Since $l^j N_j = 0$ and $m^j N_j = 0$, the equation (3.14) gives

$$(4.34) \quad D_{ik}^j N_j B_\alpha^i B_\beta^k = \frac{(\mu - 2\lambda)L}{2\tau - \rho\tau^2} M_{\alpha\beta}.$$

From (2.7), (3.8) and (4.9), we have

$$(4.35) \quad {}^*H_{\alpha\beta} - {}^*M_\alpha {}^*H_\beta = \sqrt{2\tau^2 - \rho\tau^3} (H_{\alpha\beta} + D_{jk}^i N_i B_\alpha^j B_\beta^k) - M_\alpha H_\beta.$$

Thus from (4.28), (4.34) and (4.35), we have:

Theorem 4.3. *For the Kropina change with h-vector, let the h-vector b_i be a gradient and tangential to the hypersurface F^{n-1} and satisfies the condition (4.28). Then*

1. $*F^{n-1}$ is a hyperplane of the second kind if F^{n-1} is a hyperplane of the second kind and $M_{\alpha\beta} = 0$.
2. $*F^{n-1}$ is a hyperplane of the third kind if F^{n-1} is a hyperplane of the third kind.

A Finsler space F^n is called a Landsberg space if the $(v)hv$ -torsion tensor P_{ij}^r vanishes, i.e.

$$(4.36) \quad P_{ij}^r = C_{ij|k}^r y^k = 0.$$

Taking h-covariant derivative of (1.3)(ii) and using $L_{|k} = 0 = h_{ij|k}$, (4.17), we get

$$b_{r|k} C_{ij}^r + b_r C_{ij|k}^r = 0.$$

Contracting by y^k and using (4.36), we get

$$b_{r|0} C_{ij}^r = 0,$$

which is the condition (4.28). Hence we have:

Theorem 4.4. *For the Kropina change with h-vector, let the h-vector b_i be a gradient and tangential to the hypersurface F^{n-1} of a Landsberg space F^n . Then*

1. $*F^{n-1}$ is a hyperplane of the second kind if F^{n-1} is a hyperplane of the second kind and $M_{\alpha\beta} = 0$.
2. $*F^{n-1}$ is a hyperplane of the third kind if F^{n-1} is a hyperplane of the third kind.

For the Kropina change with h -vector, let the vector b_i be parallel with respect to the Cartan connection of F^n . In view of Lemma 3.2, we have

$$(4.37) \quad *F_{jk}^i = F_{jk}^i.$$

Thus from (2.7), (4.9), (4.13) and (4.37), we have:

Theorem 4.5. *For the Kropina change with h-vector, let the vector b_i be parallel with respect to the Cartan connection of F^n and tangent to the hypersurface F^{n-1} . Then $*F^{n-1}$ is a hyperplane of the second (third) kind if and only if F^{n-1} is also a hyperplane of the second (third) kind.*

References

- [1] M. K. Gupta and P. N. Pandey, On Hypersurface of a Finsler Space with a Special Metric, *Acta Math. Hungar.*, **120** (1-2) (2008) 165-177.
- [2] M. K. Gupta and P. N. Pandey, Finsler space subjected to a Kropina change with an h -vector, arXiv:1404.6324v1 [math.DG] 25 Apr 2014.
- [3] H. Izumi, Conformal transformations of Finsler spaces II. An h -conformally flat Finsler space, *Tensor N.S.* **34** (1980), 337-359.
- [4] V. K. Kropina, On projective two-dimensional Finsler spaces with special metric, *Trudy Sem. Vector. Tensor. Anal.* **11** (1961), 277.
- [5] M. Matsumoto, The induced and intrinsic Finsler connections of a hypersurface and Finslerien projective geometry, *J. Math. Kyoto Univ.*, **25-1** (1985), 107-144.
- [6] M. Matsumoto, *Foundations of Finsler geometry and special Finsler spaces*, Kaiseisha press, Saikawa, Otsu, 520 Japan, 1986.
- [7] C. Shibata, On invariant tensors of β -changes of Finsler metrics, *J. Math. Kyoto Univ.* **24-1** 163-188.